



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Nuclear division in Ascomycetes.—GUILLIERMOND¹⁷ has continued his studies on nuclear division in the Ascomycetes, which support in all essentials the conclusions of HARPER and contravene those of MAIRE (except as to Galactinia), though they are perhaps not irreconcilable with them. However, his descriptions are not so detailed as those of HARPER in his last paper on Phylactinia, especially as it relates to the centers of spindle formation. In this paper GUILLIERMOND discusses chiefly the mother-cells of the asci and secretion. The species studied comprise *Pustularia vesiculosa*, *Aleuria cerea*, *Peziza rutilans*, *P. Catinus*, and *Galactinia succosa*.—B. M. DAVIS.

Soil waters.—CAMERON and BELL show¹⁸ that as a rule the various mineral constituents of the soil solutions exist in sufficient concentration for the growth of crops, and that the magnitude of the concentrations is *practically the same for all soils*, because, generally speaking, soils contain all the common rock forming minerals, some of each species presenting its surfaces to the solvent action of the soil water; and on account of hydrolysis of the solutes this solvent action is continuous. The paper strongly supports the previous work of the Bureau of Soils which has been so much criticised, often on a *a priori* grounds.—C. R. B.

Non-infection by rusts.—*Erysiphe graminis* has a number of biologic forms which are confined to special hosts. Thus conidia from the form on wheat will not infect barley and that on oats will not infect wheat. SALMON¹⁹ has recently shown that the reason of the non-infection is not due to inability on the part of the conidia to germinate, but because the haustoria cannot establish relations with the cells of the host plant.—B. M. DAVIS.

Endoparasitic adaptation.—SALMON²⁰ shows that *Erysiphe graminis* adapts itself readily to an endophytic life. When spores are sown on a wound in oats or barley the mycelium ramifies in the intercellular spaces and haustoria are abundantly produced. Conidiophores develop profusely and perfect conidia where they arise on a free surface; and they even break through a weak barrier when they develop in intercellular spaces.—C. R. B.

Greening of seeds.—ERNST²¹ finds that during the ripening of the fruit of *Eriobotrya japonica* the seeds become green, quite independent of light, by reason of the greening of the amyloplasts. The process begins at the plumule of the

¹⁷ GUILLIERMOND, A., Remarques sur la karyokinèse des Ascomycètes. Ann. Mycol. 3:343-361. pls. 10-12. 1905.

¹⁸ CAMERON, F. K., and BELL, J. M., The mineral constituents of soils. U. S. Dept. Agric., Bur. Soils Bull. 30. pp. 70.

¹⁹ SALMON, E. S., On the stages of development reached by certain biologic forms of *Erysiphe* in cases of non-infection. New Phytol. 4:217. 1905. pl. 5.

²⁰ SALMON, E. S., On endophytic adaptation shown by *Erysiphe graminis* DC. under cultural conditions. Phil. Trans. Roy. Soc. London B. 198:87-97. pl. 6. 1905.

²¹ ERNST, A., Das Ergrünen der Samen von *Eriobotrya japonica*. Beihefte Bot. Centralbl. 19¹: 118-130. pl. 2. 1905.